

A STUDY ON THE TRANSAMINASE ACTIVITY OF *RAILLIETINA TETRAGONA* (MOLIN, 1858) INFECTING DOMESTIC CHICK (*GALLUS DOMESTICUS*)

Achaiah N*

Department of Zoology, Kakatiya University, Warangal-506 009. (A.P), India

E-mail: achaiahsai@gmail.com

ABSTRACT

Raillietina tetragona is an endogenous helminth parasite infecting domestic chick. Transaminase enzyme (ALAT and AAT) levels in different regions representing different metabolic states were evaluated and the activity levels were statistically evaluated. A marked gradient of activity levels was observed. The results are discussed in relation to their metabolic activities.

Key words : *Raillietina tetragona*, domestic chick, Transaminases, ALAT and AAT, metabolic activities.

INTRODUCTION

Cestodes are a group of endoparasites which inhabit the intestines of the host. Much attention has been paid to carbohydrate metabolism as they are the chief energy source in cestodes. Enzymes are catalysts of biological systems. Transamination is the main mechanism for protein synthesis in helminthes (Von Brand, 1973). Transamination provides a link between carbohydrate and protein metabolism by interconverting keto acids and amino acids *vice versa*. Daugherty (1952) found the importance of carbohydrate intermediates of the Krebs cycle in protein metabolism of cestodes. These compounds, formed from glucose, apparently provide the tapeworm with the necessary building blocks for the production of amino acids and fats. Cheng (1964) stated the interrelationship between carbohydrate and protein metabolism is probably very close in certain helminthes. Yoon (1964) indicated that the transaminase plays an important role in protein synthesis in helminth parasites.

Daugherty (1952) reported high transaminase activity from *Fasciola hepatica*. Aldrich, Chandler and Daugherty (1954) studied its activity in *Hymenolepis diminuta*. Zeldon and Read (1960) evaluated the transaminase activity from three species of *Hymenolepis*. Cestodes synthesise proteins rapidly (Barrett, 1981) for egg production and tegument formation (Harris, 1983). Pathak *et al* (1981) reported serum ALAT and AAT levels of goats infected with *Cystecircus tenuicullis*. Min and Seo (1963) reported their activity from some helminthes. Nazifi *et al* (2011) and Radfar *et al* (2012) reported ALAT and AST activity levels from *Cystecircus tenuicullis*.

Although transaminases were identified in animal tissues as early as 1937, very few observations have been made on these systems in parasitic helminthes. In the present study, a comparative analysis of transaminases was conducted in different regions of *Raillietina tetragona*.

MATERIAL AND METHODS


Parasites were obtained from the naturally infected domestic chick, collected from different parts of Warangal. Collected worms were washed in saline and kept on blotting paper to remove water. Worms of same length were selected and biochemically analysed for ALAT and AAT levels. The results were analysed statistically by ANOVA.

RESULTS AND DISCUSSION

Transaminase activity levels were determined by Reitman and Frankel (1957) as modified by Bergmeyer (1965).

The results of ALAT activity levels are presented in Figure-1. ALAT activity levels in immature, mature and gravid regions are 0.216 ± 0.080 , 0.333 ± 0.087 and 0.95 ± 0.253 μ moles of sodium pyruvate /mg protein /hr.

Figure-1. Histogram showing regional distribution of ALAT activity in *R.tetragona* (Values are expressed in μ M of sodium pyruvate/mg protein/hour)




The results of AAT activity levels are presented in Figure-2. AAT activities are 0.783 ± 0.076 , 0.966 ± 0.111 and 0.733 ± 0.102 μ moles of sodium pyruvate/mg protein/irrespectively in immature, mature and gravid regions.

The present study revealed higher levels of ALAT in mature region and AAT levels in gravid region. The low activity of transaminases in *Raillitina tetragona* is in concurrence with earlier findings of Aldrich *et al* (1954) from

Hymenolepis diminuta, Wertheim *et al* (1960) from *Hymenolepis diminuta*, *Hymenolepis citelli* and *Hymenolepis nana*. When compared to vertebrates and insects, tapeworms have a limited capacity for transamination (Aldrich, 1954). It is in contrast to the higher levels of transaminases in trematodes (Von Brand, 1973). The low transaminase activity may be due to nutrient rich environment of the host.

Figure-2. Histogram showing regional distribution of AAT activity in *R.tetragona* (Values are expressed in μ M of sodium pyruvate/mg protein/hour)



The reasons for low activity may be dietary effects, age of worms and reproductive phases of worm, which could also be attributed for variation in activity among three different regions. Werethiem *et al* (1960) declared that very few compounds serve as effective amino donors in *Hymenolepis*. In comparison with vertebrates (Awapara *et al* 1952), insects (Kilby and Neville, 1957) and *Ascaris* (Savel, 1955), helminthes show limited activity (Min and Seo, 1966). They are capable of absorbing amino acids and other materials from the gut of host.

REFERENCES

1. Aldrich, D.V., A.C. Chandler, and J.W. Daugherty. (1954): Intermediary protein metabolism in helminths. II. Effect of host castration on amino acid metabolism in *Hymenolepis diminuta*. Exp. Parasit. 3: 173-184.

2. **Awapara, J. and Seale, B.** (1952): Distribution of transaminases in rat organs. *J.Biol.Chem.* 194: 497-502.
3. **Barrett, J** (1981): *Biochemistry of parasitic helminthes.* Macmillan Publishers Ltd., London.
4. **Cammarata, P.S., and Cohen, P.P** (1950): The scope of transamination reaction in animal tissues. *J.Biol.Chem.*187, 439-452.
5. **Cheng** (1974): *General Parsitology* (New York, Academic press).
6. **Campbell, J. W** (1963): Amino acids and nucleotides of the cestode, *Hymenolepis diminuta*. *Comp.Biochem. Physiol.* 8: 181-185.
7. **Conrad, F.G** (1957): *New Eng.J.Med.*256, 602.
8. **Daugherty, J. W.** (1952): Intermediary protein metabolism in helminths.I. Transaminase Reactions in *Fasciola hepatica*. *Exp. Parasit.* 1: 331-338.
9. **Daugherty** (1957): Intermediary protein metabolism in helminths. *Exp. Parasite* 6:62-67.
10. **Kilby, B. A. and Neville, E.** (1957): Amino acid metabolism in locust tissues. *J. Exp. Biol.*34: 276-289.
11. **Nizami** (1977): Quantitative studies in 7 species of digenetic trematodes
12. **Min and Seo** (1966): Studies on some transamination reactions in some helminth parasitic helminths. *Vol.4, No.2:7-13.*
13. **Reitman. S and Frankel, S** (1956): Colorimetric method for the determination of glutamic oxalo acetic and glutamic pyruvic transaminase. *Am.J.of.clin.Pathol.*28, 56-63.
14. **Smyth JD, McManus DP** (1989): *The physiology and biochemistry of cestodes.* Cambridge University Press, Cambridge.
15. **Soulsby E.J.L.** (1986): *Helmnths, Arthropods and protozoa of domesticated animals,* 7th ed. London: Bailliere and Tindall, 809 p.
16. **Werthiem.G, Zeldon.R, and Read. C.P** (1960): Transaminases of Tapeworms. *J.Parasit.*46, 496-499.
17. **Yong Ok Min and Byong Seol Seo** (1966): Studies on transaminase reactions in some parasitic helminths. *Vol.4 .No.2:7-13*
18. **Yamaguti, S.** (1961) *Systema Helminthum,* Vol.II. Cestodes of Vertebrates. New York/London, Interscience Publishers INC.